

COMPOSITE FLUXBRANES WITH GENERAL INTERSECTIONS

V.D. Ivashchuk¹,

*Center for Gravitation and Fundamental Metrology, VNIIMS, 3/1 M. Ulyanovoy Str.,
Moscow 119313, Russia*

*Institute of Gravitation and Cosmology, Peoples' Friendship University of Russia,
6 Miklukho-Maklaya Str., Moscow 117198, Russia*

Abstract

Generalized composite fluxbrane solutions for a wide class of intersection rules are obtained. The solutions are defined on a manifold which contains a product of n Ricci-flat spaces $M_1 \times \dots \times M_n$ with 1-dimensional M_1 . They are defined up to a set of functions H_s obeying non-linear differential equations equivalent to Toda-type equations with certain boundary conditions imposed. A conjecture on polynomial structure of governing functions H_s for intersections related to semisimple Lie algebras is suggested. This conjecture is valid for Lie algebras: A_m , C_{m+1} , $m \geq 1$. For simple Lie algebras the powers of polynomials coincide with the components of the dual Weyl vector in the basis of simple roots. Explicit formulas for $A_1 \oplus \dots \oplus A_1$ (orthogonal), "block-orthogonal" and A_2 solutions are obtained. Certain examples of solutions in $D = 11$ and $D = 10$ (*IIA*) supergravities (e.g. with A_2 intersection rules) and Kaluza-Klein dyonic A_2 flux tube, are considered.

*PACS numbers: 0420J, 0450, 0470

¹ivas@rgs.phys.msu.su

1 Introduction

Recently a lot of papers were devoted to multidimensional generalizations of the well-known Melvin solution [1] (for exact solutions and their applications see [3]-[32] and references therein). We remind that the original Melvin solution describes the gravitational field of a magnetic flux tube. In the works of Gibbons and Wiltshire [2] and Gibbons and Maeda [3] the Melvin solution was generalized to arbitrary dimensions (in [3] with the inclusion of dilaton), and hence the simplest fluxbranes appeared.² (A term fluxbrane was suggested in [7]. The Melvin solution is currently denoted as $F1$.) In papers [4, 5, 6, 7], devoted to Kaluza-Klein Melvin solution (e.g. non-perturbative instability and pair production of magnetically charged black holes) it was shown that $F1$ -fluxbrane supported by potential 1-form has a nice interpretation as a modding of flat space in one dimension higher. This "modding" technique is widely used in construction of new solutions in supergravitational models and also for various physical applications in string and M -theory: construction of exact string backgrounds [24, 28], duality between 0A and IIA string theories [10], dielectric effect [14, 15, 17], construction of supersymmetric configurations [18, 27] etc

The "modding" technique may be also used for Fp -fluxbranes supported by forms of higher ranks (constructed by use of 1-forms). Another approach suggested in [8, 9], which works for Fp -fluxbranes, is based on generating techniques for certain duality groups. An important result here is a construction by Chen, Gal'tsov and Sharakin [9] of intersecting $F6$ and $F3$ fluxbranes corresponding to M -branes in $D = 11$ supergravity.

The third and the most direct method is based on solving of Einstein equations [11, 12, 13]. This article is devoted to further developing of this approach using the p -brane solutions from [43] (for a review of p -brane solutions see [33, 34]). We remind, that in [43] a family of p -brane "cosmological-type" solutions with nearly arbitrary (up to some restrictions) intersection rules were obtained (see Sect. 2). These solutions are defined up to solutions to Toda-type equations and contain as a special case a subclass of solutions with cylindrical symmetry. Here we single out a subclass of generalized fluxbrane configurations related to Toda-type equations with certain asymptotical conditions imposed (Sect. 3). These fluxbrane solutions are governed by functions $H_s(z) > 0$ defined on the interval $(0, z_0)$ and obeying a set of second order non-linear differential equations

$$\frac{d}{dz} \left(\frac{z}{H_s} \frac{d}{dz} H_s \right) = \hat{B}_s \prod_{s' \in S} H_{s'}^{-A_{ss'}}, \quad (1.1)$$

with the following boundary conditions imposed:

$$(i) \quad H_s(+0) = 1, \quad (1.2)$$

$s \in S$, (S is non-empty set). In (1.1) all $\hat{B}_s \neq 0$ are constants and $(A_{ss'})$ is a "quasi-Cartan" matrix ($A_{ss} = 2$) coinciding with the Cartan one when intersections are related to Lie algebras. In most interesting examples $z_0 = +\infty$ and all $\hat{B}_s > 0$.

²The reference [2] is added in the new version

We remind that a slightly different equations occur for black brane solutions from [36, 37, 38]

$$\frac{d}{dz} \left(\frac{(1-2\mu z)}{H_s} \frac{d}{dz} H_s \right) = \bar{B}_s \prod_{s' \in S} H_{s'}^{-A_{ss'}}, \quad (1.3)$$

where $\mu > 0$ is extremality parameter, $z \in (0, (2\mu)^{-1})$ and all $\bar{B}_s \neq 0$, (in most interesting cases $\bar{B}_s < 0$). For black brane solutions the finite limits (on a horizon) $H_s((2\mu)^{-1} - 0) = H_{s0} > 0$ exist. We note that fluxbrane "master equations" (1.1) may be obtained from the black brane ones (1.3) in the limit $\mu \rightarrow \infty$. For different aspects of p-brane/fluxbrane correspondence see [16, 32].

In Sect. 4 we suggest a hypothesis: the solutions to eqs. (1.1), (1.2) are polynomials when intersection rules correspond to semisimple Lie algebras. This hypothesis (**Conjecture**) may be proved for Lie algebras A_m , C_{m+1} , $m = 1, 2, \dots$, along a line as it was done in [37, 38] for black brane solutions. It is also confirmed by special "block-orthogonal" fluxbrane solutions and explicit formulas for the solutions corresponding to Lie algebras $A_1 \oplus \dots \oplus A_1$ and A_2 . In Sect. 5 certain examples of solutions in $D = 11$ supergravity, i.e. $F6$ and $F3$ fluxbranes, $F6 \cap F3$ solutions with $A_1 \oplus A_1$ and A_2 intersection rules, $F7, (NS)F6, F5$ and (dual) $F1, (NS)F2, F3$ fluxbranes in $D = 10$ *IIA* supergravity and Kaluza-Klein dyonic A_2 -flux tube, are considered.

2 p -brane cosmological-type solutions with general intersection rules

2.1 The model

We consider a model governed by the action [35]

$$S = \int d^D x \sqrt{|g|} \left\{ R[g] - h_{\alpha\beta} g^{MN} \partial_M \varphi^\alpha \partial_N \varphi^\beta - \sum_{a \in \Delta} \frac{\theta_a}{n_a!} \exp[2\lambda_a(\varphi)] (F^a)^2 \right\} \quad (2.1)$$

where $g = g_{MN}(x) dx^M \otimes dx^N$ is a metric, $\varphi = (\varphi^\alpha) \in \mathbf{R}^l$ is a vector of scalar fields, $(h_{\alpha\beta})$ is a constant symmetric non-degenerate $l \times l$ matrix ($l \in \mathbf{N}$), $\theta_a = \pm 1$,

$$F^a = dA^a = \frac{1}{n_a!} F_{M_1 \dots M_{n_a}}^a dz^{M_1} \wedge \dots \wedge dz^{M_{n_a}} \quad (2.2)$$

is a n_a -form ($n_a \geq 1$), λ_a is a 1-form on $\mathbf{R}^l$: $\lambda_a(\varphi) = \lambda_{\alpha a} \varphi^\alpha$, $a \in \Delta$, $\alpha = 1, \dots, l$. In (2.1) we denote $|g| = |\det(g_{MN})|$,

$$(F^a)_g^2 = F_{M_1 \dots M_{n_a}}^a F_{N_1 \dots N_{n_a}}^a g^{M_1 N_1} \dots g^{M_{n_a} N_{n_a}}, \quad (2.3)$$

$a \in \Delta$. Here Δ is some finite set.

2.2 "Cosmological-type" solutions

Let us consider a family of solutions to field equations corresponding to the action (2.1) and depending upon one variable u [43]. These solutions are defined on the manifold

$$M = (u_-, u_+) \times M_1 \times M_2 \times \dots \times M_n, \quad (2.4)$$

where (u_-, u_+) is an interval belonging to $\mathbf{R}$. The solutions read [43]

$$g = \left(\prod_{s \in S} [f_s(u)]^{2d(I_s)h_s/(D-2)} \right) \left\{ \exp(2c^0 u + 2\bar{c}^0) w du \otimes du + \right. \quad (2.5)$$

$$\left. \sum_{i=1}^n \left(\prod_{s \in S} [f_s(u)]^{-2h_s \delta_{iI_s}} \right) \exp(2c^i u + 2\bar{c}^i) g^i \right\},$$

$$\exp(\varphi^\alpha) = \left(\prod_{s \in S} f_s^{h_s \chi_s \lambda_{a_s}^\alpha} \right) \exp(c^\alpha u + \bar{c}^\alpha), \quad (2.6)$$

$$F^a = \sum_{s \in S} \delta_{a_s}^a \mathcal{F}^s, \quad (2.7)$$

$\alpha = 1, \dots, l$. In (2.5) $w = \pm 1$, $g^i = g_{m_i n_i}^i(y_i) dy_i^{m_i} \otimes dy_i^{n_i}$ is a Ricci-flat metric on M_i , $i = 1, \dots, n$,

$$\delta_{iI} = \sum_{j \in I} \delta_{ij} \quad (2.8)$$

is the indicator of i belonging to I : $\delta_{iI} = 1$ for $i \in I$ and $\delta_{iI} = 0$ otherwise.

The p -brane set S is by definition

$$S = S_e \sqcup S_m, \quad S_v = \sqcup_{a \in \Delta} \{a\} \times \{v\} \times \Omega_{a,v}, \quad (2.9)$$

$v = e, m$ and $\Omega_{a,e}, \Omega_{a,m} \subset \Omega$, where $\Omega = \Omega(n)$ is the set of all non-empty subsets of $\{1, \dots, n\}$. Here and in what follows $\sqcup$ means the union of non-intersecting sets. Any p -brane index $s \in S$ has the form

$$s = (a_s, v_s, I_s), \quad (2.10)$$

where $a_s \in \Delta$ is colour index, $v_s = e, m$ is electro-magnetic index and the set $I_s \in \Omega_{a_s, v_s}$ describes the location of p -brane worldvolume.

The sets S_e and S_m define electric and magnetic p -branes correspondingly. In (2.6)

$$\chi_s = +1, -1 \quad (2.11)$$

for $s \in S_e, S_m$ respectively. In (2.7) forms

$$\mathcal{F}^s = Q_s \left(\prod_{s' \in S} f_{s'}^{-A_{ss'}} \right) du \wedge \tau(I_s), \quad (2.12)$$

$s \in S_e$, correspond to electric p -branes and forms

$$\mathcal{F}^s = Q_s \tau(\bar{I}_s), \quad (2.13)$$

$s \in S_m$, correspond to magnetic p -branes; $Q_s \neq 0$, $s \in S$. In (2.13) and in what follows

$$\bar{I} \equiv I_0 \setminus I, \quad I_0 = \{1, \dots, n\}. \quad (2.14)$$

All manifolds M_i are assumed to be oriented and connected and the volume d_i -forms

$$\tau_i \equiv \sqrt{|g^i(y_i)|} dy_i^1 \wedge \dots \wedge dy_i^{d_i}, \quad (2.15)$$

and parameters

$$\varepsilon(i) \equiv \text{sign}(\det(g_{m_i n_i}^i)) = \pm 1 \quad (2.16)$$

are well-defined for all $i = 1, \dots, n$. Here $d_i = \dim M_i$, $i = 1, \dots, n$, $D = 1 + \sum_{i=1}^n d_i$. For any $I = \{i_1, \dots, i_k\} \in \Omega$, $i_1 < \dots < i_k$, we denote

$$\tau(I) \equiv \tau_{i_1} \wedge \dots \wedge \tau_{i_k}, \quad (2.17)$$

$$M(I) \equiv M_{i_1} \times \dots \times M_{i_k}, \quad (2.18)$$

$$d(I) \equiv \dim M(I) = \sum_{i \in I} d_i, \quad (2.19)$$

$$\varepsilon(I) \equiv \varepsilon(i_1) \dots \varepsilon(i_k). \quad (2.20)$$

$M(I_s)$ is isomorphic to p -brane worldvolume (see (2.10)).

The parameters h_s appearing in the solution satisfy the relations

$$h_s = K_s^{-1}, \quad K_s = B_{ss}, \quad (2.21)$$

where

$$B_{ss'} \equiv d(I_s \cap I_{s'}) + \frac{d(I_s)d(I_{s'})}{2-D} + \chi_s \chi_{s'} \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta}, \quad (2.22)$$

$s, s' \in S$, with $(h^{\alpha\beta}) = (h_{\alpha\beta})^{-1}$. Here we assume that

$$(i) \quad B_{ss} \neq 0, \quad (2.23)$$

for all $s \in S$, and

$$(ii) \quad \det(B_{ss'}) \neq 0, \quad (2.24)$$

i.e. the matrix $(B_{ss'})$ is a non-degenerate one. In (2.12) another non-degenerate matrix (a so-called "quasi-Cartan" matrix) appears

$$(A_{ss'}) = (2B_{ss'}/B_{s's'}). \quad (2.25)$$

Here some ordering in the set S is assumed.

In (2.5), (2.6)

$$f_s = \exp(-q^s), \quad (2.26)$$

where $(q^s) = (q^s(u))$ is a solution to Toda-type equations

$$\ddot{q}^s = -B_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right), \quad (2.27)$$

with

$$B_s = \varepsilon_s K_s Q_s^2, \quad (2.28)$$

$s \in S$. Here

$$\varepsilon_s = (-\varepsilon[g])^{(1-\chi_s)/2} \varepsilon(I_s) \theta_{a_s}, \quad (2.29)$$

$s \in S$, $\varepsilon[g] \equiv \text{sign det}(g_{MN})$. More explicitly (2.29) reads: $\varepsilon_s = \varepsilon(I_s) \theta_{a_s}$ for $v_s = e$ and $\varepsilon_s = -\varepsilon[g] \varepsilon(I_s) \theta_{a_s}$, for $v_s = m$.

Vectors $c = (c^A) = (c^i, c^\alpha)$ and $\bar{c} = (\bar{c}^A)$ satisfy the linear constraints

$$U^s(c) = \sum_{i \in I_s} d_i c^i - \chi_s \lambda_{a_s \alpha} c^\alpha = 0, \quad U^s(\bar{c}) = \sum_{i \in I_s} d_i \bar{c}^i - \chi_s \lambda_{a_s \alpha} \bar{c}^\alpha = 0, \quad (2.30)$$

$s \in S$, and

$$c^0 = \sum_{j=1}^n d_j c^j, \quad \bar{c}^0 = \sum_{j=1}^n d_j \bar{c}^j, \quad (2.31)$$

$$2E = 2E_{TL} + h_{\alpha\beta} c^\alpha c^\beta + \sum_{i=1}^n d_i (c^i)^2 - \left(\sum_{i=1}^n d_i c^i \right)^2 = 0, \quad (2.32)$$

where

$$E_{TL} = \frac{1}{4} \sum_{s,s' \in S} h_s A_{ss'} \dot{q}^s \dot{q}^{s'} + \sum_{s \in S} A_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right), \quad (2.33)$$

is an integration constant (energy) for the solutions from (2.27). Here $A_s = \frac{1}{2} \varepsilon_s Q_s^2 = B_s h_s / 2$, $s \in S$. Eqs. (2.27) correspond to the Toda-type Lagrangian

$$L_{TL} = \frac{1}{4} \sum_{s,s' \in S} h_s A_{ss'} \dot{q}^s \dot{q}^{s'} - \sum_{s \in S} A_s \exp\left(\sum_{s' \in S} A_{ss'} q^{s'}\right). \quad (2.34)$$

Remark 1. Here we identify notations for g^i and $\hat{g}^i$, where $\hat{g}^i = p_i^* g^i$ is the pullback of the metric g^i to the manifold M by the canonical projection: $p_i : M \rightarrow M_i$, $i = 1, \dots, n$. An analogous agreement will be also kept for volume forms etc.

Due to (2.12) and (2.13), the dimension of p -brane worldvolume $d(I_s)$ is defined by

$$d(I_s) = n_{a_s} - 1, \quad d(I_s) = D - n_{a_s} - 1, \quad (2.35)$$

for $s \in S_e, S_m$ respectively. For a p -brane: $p = p_s = d(I_s) - 1$.

2.3 Restrictions on p -brane configurations.

The solutions presented above are valid if two restrictions on the sets of p -branes are satisfied. These restrictions guarantee the block-diagonal form of the energy-momentum tensor and the existence of the sigma-model representation (without additional constraints) [35, 34].

Let us denote $w_1 \equiv \{i | i \in \{1, \dots, n\}, d_i = 1\}$, and $n_1 = |w_1|$ (i.e. n_1 is the number of 1-dimensional spaces among M_i , $i = 1, \dots, n$).

Restriction 1. For any $a \in \Delta$ and $v = e, m$ there are no $I, J \in \Omega_{a,v}$ such that $I = \{i\} \sqcup (I \cap J)$, and $J = (I \cap J) \sqcup \{j\}$ for some $i, j \in w_1$, $i \neq j$.

Restriction 2.. For any $a \in \Delta$ there are no $I \in \Omega_{a,e}$ and $J \in \Omega_{a,m}$ such that $\bar{J} = \{i\} \sqcup I$.

Restriction 1 is satisfied for $n_1 \leq 1$ and also in the non-composite case: $|\Omega_{a,e}| + |\Omega_{a,m}| = 1$ for all $a \in \Delta$. For $n_1 \geq 2$ it forbids the following pairs of two electric or two magnetic p -branes, corresponding to the same form $F^a, a \in \Delta$:




Figure 1. A forbidden by Restriction 1 pair of two electric or two magnetic p -branes.

Here $d_i = d_j = 1$, $i \neq j$, $i, j = 1, \dots, n$. Restriction 1 may be also rewritten in terms of intersections

$$(\mathbf{R1}) \quad d(I \cap J) \leq d(I) - 2, \quad (2.36)$$

for any $I, J \in \Omega_{a,v}$, $a \in \Delta$, $v = e, m$ (here $d(I) = d(J)$).

Restriction 2 is satisfied for $n_1 = 0$. For $n_1 \geq 1$ it forbids the following electromagnetic pairs, corresponding to the same form $F^a, a \in \Delta$:




Figure 2. Forbidden by Restriction 2 electromagnetic pair of p -branes

Here $d_i = 1$, $i = 1, \dots, n$. In terms of intersections Restriction 2 reads

$$(\mathbf{R2}) \quad d(I \cap J) \neq 0, \quad (2.37)$$

for $I \in \Omega_{a,e}$ and $J \in \Omega_{a,m}$, $a \in \Delta$.

2.4 U^s -vectors and scalar products

Here we remind a minisuperspace covariant form of constraints using so called U^s -vectors [35]. The constraints (2.30) may be written in the following form

$$U^s(c) = U_A^s c^A = 0, \quad U^s(\bar{c}) = U_A^s \bar{c}^A = 0, \quad (2.38)$$

where

$$(U_A^s) = (d_i \delta_{iI_s}, -\chi_s \lambda_{\alpha a_s}), \quad (2.39)$$

$s = (a_s, v_s, I_s) \in S$, $A = (i, \alpha)$.

The quadratic constraint (2.32) reads

$$E = E_{TL} + \frac{1}{2} \hat{G}_{AB} c^A c^B = 0, \quad (2.40)$$

and

$$(\hat{G}_{AB}) = \begin{pmatrix} G_{ij} & 0 \\ 0 & h_{\alpha\beta} \end{pmatrix}, \quad (2.41)$$

is the (truncated) target space metric with

$$G_{ij} = d_i \delta_{ij} - d_i d_j, \quad (2.42)$$

$i, j = 1, \dots, n$. Let

$$(U, U') = \hat{G}^{AB} U_A U'_B, \quad (2.43)$$

be the scalar product, where $U = U_A z^A$, $U' = U'_A z^A$ are linear functions (1-forms) on $\mathbf{R}^{n+l}$, and $(\hat{G}^{AB}) = (\hat{G}_{AB})^{-1}$. The scalar products (2.43) for co-vectors U^s from (2.39) were calculated in [35]

$$(U^s, U^{s'}) = B_{ss'}, \quad (2.44)$$

$s, s' \in S$ (see (2.22)).

Intersection rules. From (2.21), (2.22) and (2.25) we get the p -brane intersection rules corresponding to the quasi-Cartan matrix $(A_{ss'})$ [41]

$$d(I_s \cap I_{s'}) = \frac{d(I_s)d(I_{s'})}{D-2} - \chi_s \chi_{s'} \lambda_{a_s} \cdot \lambda_{a_{s'}} + \frac{1}{2} K_{s'} A_{ss'}, \quad (2.45)$$

where $\lambda_{a_s} \cdot \lambda_{a_{s'}} = \lambda_{\alpha a_s} \lambda_{\beta a_{s'}} h^{\alpha\beta}$; $s, s' \in S$.

The contravariant components $U^{sA} = \hat{G}^{AB} U_B^s$ read [35]

$$U^{si} = G^{ij} U_j^s = \delta_{iI_s} - \frac{d(I_s)}{D-2}, \quad U^{s\alpha} = -\chi_s \lambda_{a_s}^\alpha, \quad (2.46)$$

$s \in S$. Here [46]

$$G^{ij} = \frac{\delta^{ij}}{d_i} + \frac{1}{2-D}, \quad (2.47)$$

$i, j = 1, \dots, n$, are the components of the matrix inverse to (G_{ij}) from (2.42). The contravariant components (2.46) appear as powers in relations for the metric and scalar fields in (2.5) and (2.6).

We note that the solution under consideration for the special case of the A_m Toda chain was obtained earlier in [42].

3 Generalized fluxbrane solutions

3.1 The choice of parameters

In what follows we put

$$w = 1, \quad d_1 = 1, \quad (3.1)$$

i.e. the manifold M_1 is one-dimensional one and

$$1 \in I_s, \quad \forall s \in S, \quad (3.2)$$

i.e. all branes contain M_1 -submanifold. We note that Restriction 2 is satisfied automatically due to (3.2).

Here we restrict ourselves to solutions with linear asymptotics at infinity

$$q^s = -\beta^s u + \bar{\beta}^s + o(1), \quad (3.3)$$

$u \rightarrow +\infty$, where $\beta^s, \bar{\beta}^s$ are constants, $s \in S$. This relation gives us an asymptotical solution to Toda type eqs. (2.27) if

$$\sum_{s' \in S} A_{ss'} \beta^{s'} > 0, \quad (3.4)$$

for all $s \in S$. In this case the energy (2.33) reads

$$E_{TL} = \frac{1}{4} \sum_{s, s' \in S} h_s A_{ss'} \beta^s \beta^{s'}. \quad (3.5)$$

We put

$$\bar{c}^A = 0, \quad \bar{\beta}^s = 0. \quad (3.6)$$

$A = (i, \alpha)$, $s \in S$. (These relations guarantee the asymptotical flatness for $u \rightarrow +\infty$ of the 2-dimensional section of the metric for $M_1 = S^1$ and $g^1 = d\phi \otimes d\phi$).

We also put

$$c^A = -\delta_1^A + \sum_{s \in S} h_s \beta^s U^{sA}, \quad (3.7)$$

$$\beta^s = 2 \sum_{s' \in S} A^{ss'}, \quad (3.8)$$

where $s \in S$, $A = (i, \alpha)$, and the matrix $(A^{ss'})$ is inverse to the matrix $(A_{ss'}) = (2(U^s, U^{s'})/(U^{s'}, U^{s'}))$. It may be verified that the constraints (2.38) and (2.40) are satisfied identically, due to conditions (3.1), (3.2).

Relations (3.7) and (3.8) look similar to analogous relations for black branes [37, 38, 34] which appear from the horizon condition. These relations play a key role in this paper. Their physical sence (possibly related to regularity of the metric at $u \rightarrow \infty$) will be clarified in a separate publication.

3.2 The main solution

We introduce a new radial variable

$$\exp(-u) = \rho \quad (3.9)$$

and denote

$$H_s = f_s e^{-\beta^s u} = e^{-q^s - \beta^s u}, \quad (3.10)$$

$s \in S$. Then the solutions (2.5)-(2.7) may be written as follows

$$g = \left(\prod_{s \in S} H_s^{2h_s d(I_s)/(D-2)} \right) \left\{ d\rho \otimes d\rho \right. \quad (3.11)$$

$$\left. + \left(\prod_{s \in S} H_s^{-2h_s} \right) \rho^2 g^1 + \sum_{i=2}^n \left(\prod_{s \in S} H_s^{-2h_s \delta_{iI_s}} \right) g^i \right\},$$

$$\exp(\varphi^\alpha) = \prod_{s \in S} H_s^{h_s \chi_s \lambda_{a_s}^\alpha}, \quad (3.12)$$

$$F^a = \sum_{s \in S} \delta_{a_s}^a \mathcal{F}^s, \quad (3.13)$$

where

$$\mathcal{F}^s = -Q_s \left(\prod_{s' \in S} H_{s'}^{-A_{ss'}} \right) \rho d\rho \wedge \tau(I_s), \quad (3.14)$$

$s \in S_e$,

$$\mathcal{F}^s = Q_s \tau(\bar{I}_s), \quad (3.15)$$

$s \in S_m$. Here $Q_s \neq 0$, $h_s = K_s^{-1}$; parameters $K_s \neq 0$ and the non-degenerate matrix $(A_{ss'})$ are defined by relations (2.21), (2.22), (2.25), $s \in S$.

Functions $H_s > 0$ obey the equations

$$\frac{d}{\rho d\rho} \left(\frac{\rho}{H_s} \frac{dH_s}{d\rho} \right) = B_s \prod_{s' \in S} H_{s'}^{-A_{ss'}}, \quad (3.16)$$

$s \in S$, where $B_s \neq 0$ are defined in (2.28). These equations follow from Toda-type equations (2.27) and the definitions (3.9) and (3.10).

It follows from (3.3), (3.6) and (3.10) that there exist finite limits

$$H_s \rightarrow 1, \quad (3.17)$$

$s \in S$, for $\rho \rightarrow 0$.

For cylindrically symmetric case

$$M_1 = S^1, \quad g^1 = d\phi \otimes d\phi, \quad (3.18)$$

$0 < \phi < 2\pi$, we get a family of composite fluxbrane solutions. They are defined up to solutions to radial equations (3.16) with the boundary conditions (3.17) imposed. In the next sections we consider several exact solutions to eqs. (3.16) and (3.17).

Another possibility

$$M_1 = \mathbf{R}, \quad g^1 = -dt \otimes dt, \quad (3.19)$$

$-\infty < t < +\infty$ (t is time variable), will lead us to generalized Milne-type solution (that may be also a subject of a separate publication).

3.3 Fluxbrane intersection rules

The fluxbrane submanifold (worldvolume) is isomorphic to $M(\bar{I}_s)$ (see (2.14), (2.18)) and has the following dimension

$$d(\bar{I}_s) = D - 1 - d(I_s), \quad (3.20)$$

$s \in S$, (see (2.35)). The fluxbrane intersection rules read

$$d(\bar{I}_s \cap \bar{I}_{s'}) = D - 1 - d(I_s) - d(I_{s'}) + d(I_s \cap I_{s'}), \quad (3.21)$$

$s \neq s'$, with p -brane intersections defined in (2.45). This relation follows from the identity

$$\bar{I}_s \cap \bar{I}_{s'} = I_0 \setminus (I_s \cup I_{s'}), \quad (3.22)$$

(see (2.14)).

We note that here we use p -brane notations for description of flux p -branes or Fp -branes. An electric (magnetic) p -brane corresponds to a magnetic (electric) $F(D-3-p)$ fluxbrane.

4 Polynomial structure of H_s for Lie algebras

Now we deal with solutions to second order non-linear differential equations (3.16) that may be rewritten as follows

$$\frac{d}{dz} \left(\frac{z}{H_s} \frac{d}{dz} H_s \right) = \frac{1}{4} B_s \prod_{s' \in S} H_{s'}^{-A_{ss'}}, \quad (4.1)$$

where $H_s(z) > 0$, $s \in S$, $z = \rho^2$. Eqs. (3.17) read

$$H_s(z = +0) = 1, \quad (4.2)$$

$s \in S$. (Here we repeat equations (1.1)-(1.2) for convenience.)

In general one may try to seek solutions of (4.1) in a class of functions analytical in some disc $|z| < L$:

$$H_s(z) = 1 + \sum_{k=1}^{\infty} P_s^{(k)} z^k, \quad (4.3)$$

where $P_s^{(k)}$ are constants, $s \in S$. Substitution of (4.3) into (4.1) gives us an infinite chain of relations on parameters $P_s^{(k)}$ and B_s . The first relation in this chain

$$P_s \equiv P_s^{(1)} = \frac{1}{4} B_s, \quad (4.4)$$

$s \in S$, corresponds to the z^0 -term in the decomposition of (4.1). For analytical function $H_s(z)$ (4.3) ($z = \rho^2$) the metric (3.11) is regular at $\rho = 0$.

4.1 Solutions for Lie algebra $A_1 \oplus \dots \oplus A_1$

4.1.1 Orthogonal case

Meanwhile there exist solutions to eqs. (4.1)-(4.2) of polynomial type. The simplest example occurs in orthogonal case, when

$$(U^s, U^{s'}) = B_{ss'} = 0, \quad (4.5)$$

for $s \neq s'$, $s, s' \in S$. In this case $(A_{ss'}) = \text{diag}(2, \dots, 2)$ is a Cartan matrix for semisimple Lie algebra $A_1 \oplus \dots \oplus A_1$ and

$$H_s(z) = 1 + P_s z, \quad (4.6)$$

with $P_s \neq 0$ satisfying (4.4). (For A_1 -case see [13]).

4.1.2 Block-orthogonal case

The solution (4.6) may be generalized to so-called "block-orthogonal" (BO) case:

$$S = S_1 \sqcup \dots \sqcup S_k, \quad (4.7)$$

$S_i \neq \emptyset$, i.e. the set S is a union of k non-intersecting (non-empty) subsets $S_1, \dots, S_k$, and

$$(U^s, U^{s'}) = 0 \quad (4.8)$$

for all $s \in S_i$, $s' \in S_j$, $i \neq j$; $i, j = 1, \dots, k$. (For "block-orthogonal" black branes see [39, 40].) In this case (4.6) is modified as follows

$$H_s(z) = (1 + \bar{P}_s z)^{\beta^s}, \quad (4.9)$$

where β^s are defined in (3.8) and parameters $\bar{P}_s$ are coinciding inside blocks, i.e. $\bar{P}_s = \bar{P}_{s'}$ for $s, s' \in S_i$, $i = 1, \dots, k$, and satisfy the following relations

$$\bar{P}_s = B_s / (4\beta^s), \quad (4.10)$$

for $\beta^s \neq 0$, $s \in S$. In this case H_s are analytical in $|z| < L$, where $L = \min(|\bar{P}_s|^{-1}, s \in S)$.

Let $(A_{ss'})$ be a Cartan matrix for a finite-dimensional semisimple Lie algebra $\mathcal{G}$. In this case all powers in (3.8) are natural numbers

$$\beta^s = 2 \sum_{s' \in S} A^{ss'} = n_s \in \mathbf{N}, \quad (4.11)$$

and hence, all functions H_s are polynomials, $s \in S$. Integers n_s coincide with the components of the twice dual Weyl vector in the basis of simple roots [44].

Conjecture. *Let $(A_{ss'})$ be a Cartan matrix for a semisimple finite-dimensional Lie algebra $\mathcal{G}$. Then the solution to eqs. (4.1), (4.2) (if exists) is a polynomial*

$$H_s(z) = 1 + \sum_{k=1}^{n_s} P_s^{(k)} z^k, \quad (4.12)$$

where $P_s^{(k)}$ are constants, $k = 1, \dots, n_s$, integers n_s are defined in (4.11) and $P_s^{(n_s)} \neq 0$, $s \in S$.

This conjecture may be verified for Lie algebras A_m , C_{m+1} repeating all arguments from [37, 38] with the replacement of $F(z) = 1 - 2\mu z$ by $F(z) = z$.

4.2 Solutions for Lie algebra A_2

Let us consider the Lie algebra $A_2 = sl(3)$ with the Cartan matrix

$$(A_{ss'}) = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} \quad (4.13)$$

According to the **Conjecture** we seek the solutions to eqs. (4.1)-(4.2) in the following form ($n_1 = n_2 = 2$):

$$H_s = 1 + P_s z + P_s^{(2)} z^2, \quad (4.14)$$

where P_s and $P_s^{(2)} \neq 0$ are constants, $s = 1, 2$.

The substitution of (4.14) into equations (4.1) and decomposition in powers of z lead us to relations (4.4) and

$$P_s^{(2)} = \frac{1}{4}P_1P_2. \quad (4.15)$$

Thus, in the A_2 -case the solution is described by relations (3.11)-(3.15) with $S = \{s_1, s_2\}$, p -brane intersection rules (2.45), or, equivalently,

$$d(I_{s_1} \cap I_{s_2}) = \frac{d(I_{s_1})d(I_{s_2})}{D-2} - \chi_{s_1}\chi_{s_2}\lambda_{a_{s_1}} \cdot \lambda_{a_{s_2}} - \frac{1}{2}K, \quad (4.16)$$

$$d(I_{s_i}) - \frac{(d(I_{s_i}))^2}{D-2} + \lambda_{a_{s_i}} \cdot \lambda_{a_{s_i}} = K, \quad (4.17)$$

where $K = K_{s_i} \neq 0$, and functions $H_{s_i} = H_i$ are defined by relations (4.14), (4.4), (4.15) with $z = \rho^2$, $i = 1, 2$.

5 Examples of fluxbrane solutions

Here we present certain examples of fluxbrane solutions with $M_1 = S^1$ and $g^1 = d\phi \otimes d\phi$. In all examples below the total metric g has the signature $(-, +, \dots, +)$ and all (p -brane) signature parameters are positive: $\varepsilon_s = \varepsilon(I_s) = +1$ (here all $\theta_a = 1$). In what follows $0 < \rho < +\infty$. In all examples the metrics are regular at $\rho = 0$.

5.1 Solutions for algebra A_1

We start with single fluxbrane solutions ($S = \{s\}$).

5.1.1 Melvin solution ($F1$ fluxbrane)

Let $D = 4$, $n = 3$, $M_2 = \mathbf{R}$ with $g^2 = -dt \otimes dt$ and $M_3 = \mathbf{R}$ with $g^3 = d\eta \otimes d\eta$, and $I_s = \{1\}$. The solution reads [1]

$$g = H^2 \left\{ d\rho \otimes d\rho + H^{-4} \rho^2 d\phi \otimes d\phi - dt \otimes dt + d\eta \otimes d\eta \right\}. \quad (5.1)$$

$$F = -QH^{-2} \rho d\rho \wedge d\phi, \quad (5.2)$$

where $H = 1 + \frac{1}{8}Q^2\rho^2$. Here $-Q$ is proportional to magnetic field in the core.

5.1.2 $F6$ fluxbrane (corresponding to $M2$ -brane)

Consider $D = 11$ supergravity with the metric g and 4-form F in the bosonic sector [45]. Let $n = 3$, M_3 be 7-dimensional (Ricci-flat) manifold with the metric $g^3 = g_{\mu\nu}^3 dx^\mu \otimes dx^\nu$ of signature $(-, +, \dots, +)$ and M_2 be 2-dimensional (flat) manifold of signature $(+, +)$ with the metric $g^2 = g_{mn}^2 dy^m \otimes dy^n$ and $I_s = \{1, 2\}$. The solution reads

$$g = H^{1/3} \left\{ d\rho \otimes d\rho + H^{-1}(\rho^2 d\phi \otimes d\phi + g^2) + g^3 \right\}, \quad (5.3)$$

$$F = -QH^{-2} \rho d\rho \wedge d\phi \wedge \tau_2, \quad (5.4)$$

where $H = 1 + \frac{1}{2}Q^2\rho^2$. For flat g^3 this solution was obtained earlier in [9].

5.1.3 $F3$ fluxbrane (corresponding to $M5$ -brane)

Now we consider the solution dual to $F6$. Let $n = 3$, M_3 be 4-dimensional (Ricci-flat) manifold with the metric $g^3 = g_{\mu\nu}^3 dx^\mu \otimes dx^\nu$ of signature $(-, +, +, +)$, and M_2 be 5-dimensional (Ricci-flat) manifold of signature $(+, \dots, +)$ with the metric $g^2 = g_{mn}^2 dy^m \otimes dy^n$ and $I_s = \{1, 2\}$. The solution reads

$$g = H^{2/3} \left\{ d\rho \otimes d\rho + H^{-1}(\rho^2 d\phi \otimes d\phi + g^2) + g^3 \right\}, \quad (5.5)$$

$$F = Q\tau_3, \quad (5.6)$$

where $H = 1 + \frac{1}{2}Q^2\rho^2$. For flat g^2 and g^3 see [9].

5.1.4 $F7, F6, F5$ and $F1, F2, F3$ fluxbranes in IIA supergravity

The bosonic part of action for $D = 10$ IIA supergravity reads

$$S = \int d^{10}z \sqrt{|g|} \left\{ R[g] - (\partial\varphi)^2 - \sum_{a=2}^4 e^{2\lambda_a\varphi} (F^a)^2 \right\} - \frac{1}{2} \int F^4 \wedge F^4 \wedge A^2, \quad (5.7)$$

where $F^a = dA^{a-1} + \delta_4^a A^1 \wedge F^3$ is an a -form, $a = 2, 3, 4$, and

$$\lambda_2 = 3\lambda_4, \quad \lambda_3 = -2\lambda_4, \quad \lambda_4^2 = 1/8. \quad (5.8)$$

The dimensions of p -brane worldvolumes are

$$d = d(a, \chi) = \begin{cases} 1, 2, 3 & \text{in electric case, } \chi = +1, \\ 7, 6, 5 & \text{in magnetic case, } \chi = -1, \end{cases} \quad (5.9)$$

for $a = 2, 3, 4$, respectively.

The fluxbrane solutions read

$$g = H^{d/8} \left\{ d\rho \otimes d\rho + H^{-1}\rho^2 d\phi \otimes d\phi + H^{-1}g^2 + g^3 \right\}, \quad (5.10)$$

$$\exp(2\varphi) = H^{\chi\lambda_a}, \quad (5.11)$$

and

$$F^a = -QH^{-2}\rho d\rho \wedge \tau_2, \quad \text{for } \chi = +1, \quad (5.12)$$

or

$$F^a = Q\tau_3, \quad \text{for } \chi = -1, \quad (5.13)$$

with $H = 1 + \frac{1}{2}Q^2\rho^2$, $a = 2, 3, 4$.

For $\chi = +1$ we get a $F(9-a)$ fluxbrane with $d_2 = a-2$, $d_3 = 10-a$, and for $\chi = -1$ we obtain a (dual) $F(a-1)$ fluxbrane with $d_2 = 8-a$, $d_3 = a$, ($a = 2, 3, 4$). (Here the presence of Chern-Simons terms does not modify the solutions from Sect. 3.2.)

5.2 Solution for algebra $A_1 \oplus A_1$

$F6 \cap F3$ fluxbranes. We put $n = 5$, $d_2 = 1$, $g^2 = dy^2 \otimes dy^2$, $d_3 = 1$, $g^3 = dy^3 \otimes dy^3$, $d_4 = 4$ and g^4 has the Euclidean signature, $d_5 = 3$ and g^5 has the signature $(-, +, +)$, $I_{s_e} = \{1, 2, 3\}$ and $I_{s_m} = \{1, 2, 4\}$. The solution has the following form

$$g = H_e^{1/3} H_m^{2/3} \left\{ d\rho \otimes d\rho + H_e^{-1} H_m^{-1} (\rho^2 d\phi \otimes d\phi + g^2) + H_e^{-1} g^3 + H_m^{-1} g^4 + g^5 \right\}, \quad (5.14)$$

$$F = -Q_e H_e^{-2} \rho d\rho \wedge d\phi \wedge dy_2 \wedge dy_3 + Q_m dy_3 \wedge \tau_5, \quad (5.15)$$

where $H_s = 1 + \frac{1}{2} Q_s^2 \rho^2$, $s = e, m$. For flat g^i see [9].

5.3 Solutions for algebra A_2

5.3.1 $F6 \cap F3$ fluxbranes with A_2 -intersection

Now we consider a new $F6 \cap F3$ fluxbrane configuration with (a non-standard) A_2 intersection rules defined on the manifold

$$M = (0, +\infty) \times M_1 \times M_2 \times M_3 \times M_4, \quad (5.16)$$

where $d_2 = 2$, $d_3 = 5$, $d_4 = 2$. The solution is following

$$g = H_e^{1/3} H_m^{2/3} \left\{ d\rho \otimes d\rho + H_e^{-1} H_m^{-1} \rho^2 d\phi \otimes d\phi + H_e^{-1} g^2 + H_m^{-1} g^3 + g^4 \right\}, \quad (5.17)$$

$$F = -Q_e H_e^{-2} H_m \rho d\rho \wedge d\phi \wedge \tau_2 + Q_m \tau_2 \wedge \tau_4, \quad (5.18)$$

where metrics g^2 and g^3 are (Ricci-flat) metrics of Euclidean signature, g^4 is the (flat) metric of the signature $(-, +)$ and

$$H_s = 1 + P_s \rho^2 + \frac{1}{4} P_1 P_2 \rho^4, \quad (5.19)$$

where $P_s = \frac{1}{2} Q_s^2$, $s = e, m$.

5.3.2 Dyonic flux tube in Kaluza-Klein model

Let us consider 4-dimensional model

$$S = \int_M d^4 z \sqrt{|g|} \left\{ R[g] - g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi - \frac{1}{2!} \exp[2\lambda\varphi] F^2 \right\} \quad (5.20)$$

with scalar field φ , two-form $F = dA$ and $\lambda = -\sqrt{3/2}$. This model originates after Kaluza-Klein reduction of 5-dimensional gravity. The 5-dimensional metric in this case reads

$$g^{(5)} = \Phi g_{\mu\nu} dx^\mu \otimes dx^\nu + \Phi^{-2} (dy + \mathcal{A}) \otimes (dy + \mathcal{A}), \quad (5.21)$$

where

$$\mathcal{A} = \sqrt{2} A = \sqrt{2} A_\mu dx^\mu, \quad \Phi = \exp(2\varphi/\sqrt{6}). \quad (5.22)$$

We consider "dyonic" flux-tube solution defined on the manifold

$$M = (0, +\infty) \times (M_1 = S^1) \times M_2, \quad (5.23)$$

where $M_2 = \mathbf{R}^2$ and $g^2 = -dt \otimes dt + d\eta \otimes d\eta$. This solution reads

$$g = (H_e H_m)^{1/2} \left\{ d\rho \otimes d\rho - H_e^{-1} H_m^{-1} \rho^2 d\phi \otimes d\phi - dt \otimes dt + d\eta \otimes d\eta \right\}, \quad (5.24)$$

$$\exp \varphi = H_e^{\lambda/2} H_m^{-\lambda/2}, \quad (5.25)$$

$$F = dA = -Q_e H_e^{-2} H_m \rho d\rho \wedge d\phi + Q_m dt \wedge d\eta, \quad (5.26)$$

where functions H_s are defined by relations (5.19),

For 5-metric we obtain from (5.21)-(5.25)

$$g^{(5)} = H_m \{ d\rho \otimes d\rho + H_e^{-1} H_m^{-1} \rho^2 d\phi \otimes d\phi - dt \otimes dt + d\eta \otimes d\eta \} \\ + H_e H_m^{-1} (dy + \mathcal{A}) \otimes (dy + \mathcal{A}), \quad (5.27)$$

$d\mathcal{A} = \sqrt{2}F$. For $Q_m \rightarrow 0$ we get the solution from [3] and for $Q_e \rightarrow 0$ we are led to its dual version (see [9]).

6 Conclusions

Thus, here we obtained a family of fluxbrane solutions with general intersection rules. (See relations (2.23), (2.24) and Restriction 1. Restriction 2 is satisfied, since all p -branes have a common M_1 .) These solutions are given by relations (3.11)-(3.17). The metrics of solutions contain n Ricci-flat metrics. The solutions are defined up to a set of "moduli" functions H_s obeying a set of equations (3.16) with the boundary conditions (3.17) imposed. These solutions are new and generalize a lot of special fluxbrane solutions considered earlier in the literature.

Here we suggested a conjecture on polynomial structure of H_s for intersections related to semisimple Lie algebras. This conjecture is valid for Lie algebras A_m and C_{m+1} , $m \geq 1$, at may be verified with a little modification of the proof for the black brane case [37, 38].

We obtained explicit formulas for $A_1 \oplus \dots \oplus A_1$ (orthogonal), block-orthogonal and A_2 solutions. These formulas are illustrated by certain examples of solutions in $D = 10, 11$ supergravities (e.g. with A_2 intersection rules) and Kaluza-Klein dyonic A_2 flux tube. Explicit relations for H_s corresponding to other examples of Lie algebras (e.g. A_3 , B_2 etc) will be a subject of further publications. (Another topic of interest is related to supersymmetric composite fluxbranes on product of Ricci-flat manifolds. For M -branes see [47] and references therein).

Acknowledgments

This work was supported in part by the Russian Ministry for Science and Technology, Russian Foundation for Basic Research, DFG foundation and project SEE.

References

- [1] M.A. Melvin, Pure magnetic and electric geons, *Phys. Lett.* **8**, 65 (1964).
- [2] G.W. Gibbons and D.L. Wiltshire, Spacetime as a membrane in higher dimensions, *Nucl. Phys.* **B 287**, 717-742 (1987); hep-th/0109093.
- [3] G. Gibbons and K. Maeda, Black holes and membranes in higher dimensional theories with dilaton fields, *Nucl. Phys.* **B 298**, 741-775 (1988).
- [4] F. Dowker, J.P. Gauntlett, D.A. Kastor and J. Traschen, Pair creation of dilaton black holes, *Phys. Rev.* **D 49**, 2909-2917 (1994); hep-th/9309075.
- [5] F. Dowker, J.P. Gauntlett, S.B. Giddings and G.T. Horowitz, On pair creation of extremal black holes and Kaluza-Klein monopoles, *Phys. Rev.* **D 50**, 2662 (1994); hep-th/9312172.
- [6] F. Dowker, J.P. Gauntlett, G.W. Gibbons and G.T. Horowitz, The decay of magnetic fields in Kaluza-Klein theory, *Phys. Rev.* **D 52**, 6929 (1995) hep-th/9507143.
- [7] H.F. Dowker, J.P. Gauntlett, G.W. Gibbons and G.T. Horowitz, Nucleation of P -branes and fundamental strings, *Phys. Rev.* **D 53**, 7115 (1996); hep-th/9512154.
- [8] D.V. Gal'tsov and O.A. Rytchkov, Generating branes via sigma models, *Phys. Rev.* **D 58**, 122001 (1998); hep-th/9801180.
- [9] C.-M. Chen, D.V. Gal'tsov and S.A. Sharakin, Intersecting M -fluxbranes, *Grav. Cosmol.* **5**, 1(17), 45-48 (1999); hep-th/9908132.
- [10] M.S. Costa and M. Gutperle, The Kaluza-Klein Melvin solution in M-theory, *JHEP* **0103**, 027 (2001); hep-th/0012072.
- [11] P.M. Saffin, Gravitating fluxbranes, *Phys. Rev.* **D 64**, 024014 (2001); gr-qc/0104014.
- [12] M. Gutperle and A. Strominger, Fluxbranes in string theory, *JHEP* **0106**, 035 (2001); hep-th/0104136.
- [13] C.M. Chen, D.V. Gal'tsov and P.M. Saffin, Supergravity fluxbranes in various dimensions, hep-th/0110164.
- [14] M.S. Costa, C.A. Herdeiro and L. Cornalba, Flux-branes and the dielectric effect in string theory, hep-th/0105023.
- [15] R. Emparan, Tubular branes in fluxbranes, *Nucl. Phys.* **B 610**, 169 (2001) hep-th/0105062.
- [16] P.M. Saffin, Fluxbranes from p-branes, *Phys. Rev.* **D 64**, 104008 (2001); hep-th/0105220.
- [17] D. Brecher and P.M. Saffin, A note on the supergravity description of dielectric branes, *Nucl. Phys.* **B 613**, 218 (2001), hep-th/0106206.
- [18] J. Figueroa-O'Farrill and J. Simon, Generalized supersymmetric fluxbranes, hep-th/0110170.
- [19] L. Motl, Melvin matrix models, hep-th/0107002.
- [20] T. Suyama, Melvin background in heterotic theories, hep-th/0107116.

- [21] A.M. Uranga, Wrapped fluxbranes, hep-th/0108196.
- [22] T. Suyama, Properties of string theory on Kaluza-Klein Melvin background, hep-th/0110077.
- [23] T. Takayanagi and T. Uesugi, Orbifolds as Melvin geometry, hep-th/0110099.
- [24] J.G. Russo and A.A. Tseytlin, Exactly solvable string models of curved space-time backgrounds, *Nucl. Phys.* **B 449**, 91 (1995); hep-th/9502038.
- [25] O. Bergman and M.R. Gaberdiel, Dualities of type 0 strings, *JHEP* **9907**, 022 (1999) hep-th/9906055.
- [26] J.G. Russo and A.A. Tseytlin, Magnetic backgrounds and tachyonic instabilities in closed superstring theory and M-theory, *Nucl. Phys.* **B 611**, 93 (2001) hep-th/0104238.
- [27] J.G. Russo and A.A. Tseytlin, Supersymmetric fluxbrane intersections and closed string tachyons, hep-th/0110107.
- [28] A.A. Tseytlin, Magnetic backgrounds and tachyonic instabilities in closed string theory, hep-th/0108140.
- [29] E. Dudas and J. Mourad, D-branes in string theory Melvin backgrounds, hep-th/0110186.
- [30] T. Takayanagi and T. Uesugi, D-branes in Melvin background, hep-th/0110200.
- [31] J.M. Figueroa-O'Farrill and G. Papadopoulos, Homogeneous fluxes, branes and a maximally supersymmetric solution of M -theory, *JHEP* **0106**, 036 (2001); hep-th/0105308.
- [32] R. Emparan and M. Gutperler, From p-branes to fluxbranes and back, hep-th/0111177.
- [33] K.S. Stelle, Lectures on supergravity p-branes, hep-th/9701088.
- [34] V.D. Ivashchuk and V.N. Melnikov, Exact solutions in multidimensional gravity with antisymmetric forms, topical review, *Class. Quantum Grav.*, **18** (2001) R87-R152; hep-th/0110274.
- [35] V.D. Ivashchuk and V.N. Melnikov, Sigma-model for the generalized composite p-branes, hep-th/9705036; *Class. Quantum Grav.* **14**, 3001-3029 (1997); Corrigenda *ibid.* **15** (12), 3941 (1998).
- [36] V.D. Ivashchuk and V.N. Melnikov, P-brane black holes for general intersections, *Grav. Cosmol.* **5**, No 4 (20), 313-318 (1999); gr-qc/0002085.
- [37] V.D. Ivashchuk and V.N. Melnikov, Black hole p-brane solutions for general intersection rules, *Grav. Cosmol.* **6**, No 1 (21), 27-40 (2000); hep-th/9910041.
- [38] V.D. Ivashchuk and V.N. Melnikov, Toda p-brane black holes and polynomials related to Lie algebras. *Class. Quantum Grav.* **17**, 2073-2092 (2000); math-ph/0002048.
- [39] K.A. Bronnikov, Block-orthogonal brane systems, black holes and wormholes, hep-th/9710207; *Grav. Cosmol.* **4**, No 1 (13), 49 (1998).
- [40] S. Cotsakis, V.D. Ivashchuk and V.N. Melnikov, P-branes black holes and post-Newtonian approximation, *Grav. Cosmol.* **5**, No 1 (17), 52-57 (1999); gr-qc/9902148.

- [41] V.D. Ivashchuk and V.N. Melnikov, Cosmological and spherically symmetric solutions with intersecting p-branes. *J. Math. Phys.*, 1999, **40** (12), 6558-6576.
- [42] V.R. Gavrilov and V.N. Melnikov, Toda chains associated with Lie algebras A_m in multidimensional gravitation and cosmology with intersecting p -branes, *Theor. Math. Phys.* **123**, No 3, 374-394 (2000) (in Russian).
- [43] V.D. Ivashchuk and S.-W. Kim, Solutions with intersecting p-branes related to Toda chains, *J. Math. Phys.* **41**, 444-460 (2000); hep-th/9907019.
- [44] J. Fuchs and C. Schweigert, Symmetries, Lie algebras and Representations. A graduate course for physicists (Cambridge University Press, Cambridge, 1997).
- [45] E. Cremmer, B. Julia, J. Scherk. *Phys. Lett.* **B 76**, 409 (1978).
- [46] V.D. Ivashchuk, V.N. Melnikov and A.I. Zhuk, *Nuovo Cimento* **B 104**, 575 (1989).
- [47] V.D. Ivashchuk, On supersymmetric solutions in $D = 11$ supergravity on product of Ricci-flat spaces, hep-th/0012263; *Grav. Cosmol.* **6**, No. 4(24), 344-350 (2000); hep-th/0012263.